


SAVE A LIFE. GET NALOXONE.

Naloxone stops an overdose caused by opioid pain medication, methadone or heroin.

People at risk for overdose and their family and friends can learn to spot an overdose and respond to save a life.

To get naloxone, present this card to the pharmacy staff.

A



MULTI-STEP NASAL SPRAY

DIRECTIONS: Spray 1 mL (half of the syringe) into each nostril.

NO BRAND NAME/GENERIC

COST: \$-\$\$

B



SINGLE-STEP NASAL SPRAY

DIRECTIONS: Spray full dose into one nostril.

BRAND NAME: Narcan

COST: \$\$\$

C



INTRAMUSCULAR INJECTION

DIRECTIONS: Inject 1 mL in shoulder or thigh.

NO BRAND NAME/GENERIC

COST: \$-\$\$

D



AUTO-INJECTOR

DIRECTIONS: Use as directed by voice-prompt. Press black side firmly on outer thigh.

BRAND NAME: Evzio

COST: \$\$\$\$*

*Coupons available, see evzio.com for more info

FOR ALL PRODUCTS, repeat naloxone administration after 2–3 minutes if there is no response.

Most insurance will cover at least one of these options, or you can pay cash. All products contain at least two doses

For more on opioid safety, videos on how to use naloxone, or to get help for addiction, go to PrescribetoPrevent.org

TECHNICIAN: Use product information on back of card to fill and process insurance or provide cash price to patient. Get pharmacist for verification and counseling.

PHARMACIST: Counsel patient on overdose prevention and the **steps of overdose response**.

STEPS OF OVERDOSE RESPONSE:

1. Check for response.
2. Call 911.
3. Give naloxone. If no response in 2-3 minutes, repeat dose.
4. Give rescue breaths or chest compressions.
5. Stay with person until help arrives.

A



MULTI-STEP NASAL SPRAY

DIRECTIONS: Spray 1 mL (half of the syringe) into each nostril.

QUANTITY: #2 x2 mL Luer-Jet™ Luer-Lock needleless syringe plus #2 mucosal atomizer devices (MAD-300)

SIG: Spray 1 mL (1/2 of syringe) into each nostril.

NDC: 76329-3369-01

COST: \$-\$\$

B



SINGLE-STEP NASAL SPRAY

DIRECTIONS: Spray full dose into one nostril.

QUANTITY: #1 two-pack of two 4mg/0.1 mL intranasal devices)

SIG: Spray .1 mL into one nostril.

NDC: 69547-353-02

COST: \$\$\$

C



INTRAMUSCULAR INJECTION

DIRECTIONS: Inject 1 mL in shoulder or thigh.

QUANTITY: #2 single-use 1 mL vials PLUS #2 3 mL syringe w/ 23-25 gauge 1-1.5 inch IM needles

SIG: Inject 1 mL in shoulder or thigh.

NDC: 00409-1215-01
67457-0292-02

COST: \$-\$\$

D



AUTO-INJECTOR

DIRECTIONS: Use as directed by voice-prompt. Press black side firmly on outer thigh.

QUANTITY: #1 two-pack of two 0.4 mg/0.4 mL prefilled auto-injector devices

SIG: Inject into outer thigh as directed by English voice-prompt system. Place black side firmly on outer thigh and depress and hold for 5 seconds.

NDC: 60842-030-01

COST: \$\$\$\$*

*Coupons available, see evzio.com for more info

Naloxone stops an overdose caused by opioid pain medication, methadone or heroin.

People at risk for overdose and their family and friends can learn to spot an overdose and respond to save a life.

For more on opioid safety, videos on how to use naloxone, or to get help for addiction, go to PrescribetoPrevent.org